



The American College of
Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

Physician Script on Influenza Immunization During Pregnancy

All women should receive the influenza vaccine; this is particularly important during pregnancy and the postpartum period. The influenza vaccination is an essential element of prenatal care because pregnant women are at an increased risk of serious illness and mortality due to influenza. In addition, maternal vaccination is the most effective strategy to protect newborns because the vaccine is not approved for use in infants younger than 6 months.

- Only the inactivated influenza vaccine is recommended during pregnancy. Live, attenuated influenza vaccine, which is given as a nasal spray, is contraindicated for pregnant women.
- Inadvertent administration of the live, attenuated influenza vaccine during pregnancy has not been shown to be harmful. The live, attenuated influenza vaccine is safe to administer postpartum and to family members.
- It is safe for pregnant women to receive a vaccine with thimerosal. Thimerosal, a mercury-containing preservative used in multidose vials, has not been shown to cause any adverse effects except for occasional local skin reactions.
- There is no scientific evidence that thimerosal-containing vaccines cause adverse effects in children born to women who received vaccines with thimerosal.
- Some women may still be concerned; preservative free single-dose influenza vaccines are available from certain manufacturers.
- It is required by law for you to give your patients a Vaccine Information Statement upon receiving an immunization. Vaccine Information Statement forms can be found in multiple languages at www.immunize.org/vis/.
- Multiple studies show that the most effective way to increase your patient's vaccination acceptance rate is for you to directly recommend and provide the vaccine. Talk to your patients about the flu shot today. Here is a script for your consideration:

"I strongly recommend you get the flu shot today. I offer the influenza vaccine to all of my pregnant patients and to women who are considering becoming pregnant. The vaccine is safe and effective for pregnant women. The risks of getting sick with the flu are far greater for a pregnant woman and her baby than the possibility of having a complication from the vaccine. The flu shot will protect you as well as your baby in the first 6 months of life from getting the flu. Your family members who have contact with your newborn also should be vaccinated."

RESOURCES

The American College of Obstetricians and Gynecologists. Seasonal influenza (flu) for ob-gyns. Available at: http://www.immunizationforwomen.org/immunization_facts/seasonal_influenza. Retrieved April 1, 2013.

Centers for Disease Control and Prevention. Pregnant women and influenza (flu). Available at: <http://www.cdc.gov/flu/protect/vaccine/pregnant.htm>. Retrieved April 1, 2013.

Department of Health and Human Services. Pregnant women at risk for the flu. Available at: <http://www.flu.gov/at-risk/pregnant/index.html>. Retrieved April 1, 2013



Coding Information on Influenza Immunization for Pregnant Patients

CPT Codes for Vaccine Administration

Code	Method	Route of Administration	Type of Service	Reporting Rules
90471	Injection	Percutaneous, intradermal, subcutaneous, or intramuscular	Primary	Report only one primary vaccine administration per encounter.
+90472	Injection	Percutaneous, intradermal, subcutaneous, or intramuscular	Additional	Report for secondary or subsequent vaccine administration. Report only with code 90471.
90460	Any route	Percutaneous, intradermal, subcutaneous, or intramuscular	Primary	Report only one primary vaccine administration per day. Report for administration of first vaccine if more than one was provided. Physician also provides counseling. Patient is 18 years of age or younger.
90461	Any route	Percutaneous, intradermal, subcutaneous, or intramuscular	Additional	Report for secondary or subsequent vaccine administration. Physician also provides counseling. Patient is 18 years of age or younger.

Influenza Vaccines Administered to Adolescents and Adults

Vaccine	Code for Vaccine Product	CPT Administration Code
Influenza virus, split, preservative free, patients 3 years of age and older, intramuscular	90656	90471–90472
Influenza virus, split, patients 3 years of age and older, intramuscular	90658	90471–90472

For more information, please visit the Coding section on the Immunization for Women web page, http://www.immunizationforwomen.org/practice_management/coding.

This guide is provided by the American College of Obstetricians and Gynecologists (the College) for educational purposes only. It is not intended to represent the only, or necessarily the best, coding format or method for the situations discussed, but rather as an approach, view, statement, or opinion that may be helpful to persons responsible for diagnosis and procedure coding. The statements made in this publication should not be construed as the College policy or procedure, nor as standards of care. The American College of Obstetricians and Gynecologists makes no representations or warranties, expressed or implied, regarding the accuracy of the information contained in this document and disclaims any liability or responsibility for any consequences resulting from or otherwise related to any use of, or reliance on, this resource.