

Screening Pregnant Women for Hepatitis B Virus (HBV) Infection:

Ordering Prenatal Hepatitis B Surface Antigen (HBsAg) Tests from Major Commercial Laboratories

| Laboratory | Test Option | Test Name | Reflex to Confirmation Test* | Test Code/ID | CPT Code | Web Link |
|---------------------------|-------------|---|------------------------------|--------------|----------|---|
| ARUP Laboratories | Panel | Prenatal Reflexive Panel | ✓ | 0095044 | 87340** | http://ltd.aruplab.com/Tests/Pub/0095044 |
| | Standalone | Hepatitis B Virus Surface Antigen with Reflex to Confirmation, Prenatal | ✓ | 2007573 | 87340 | http://ltd.aruplab.com/Tests/Pub/2007573 |
| LabCorp | Panel | Prenatal Profile I with Hepatitis B Surface Antigen | ✓ | 202945 | 80055 | https://www.labcorp.com/wps/portal/provider/testmenu/ (Enter test code or CPT code to search for test) |
| | Panel | Hepatitis Profile XIII (HBV Prenatal Profile) | ✓ | 265397 | 87340** | https://www.labcorp.com/wps/portal/provider/testmenu/ (Enter test code or CPT code to search for test) |
| | Standalone | N/A | N/A | N/A | N/A | |
| Mayo Medical Laboratories | Panel | Prenatal Hepatitis Evaluation | ✓ | PHSP | 87340** | http://www.mayomedicallaboratories.com/test-catalog/Overview/5566 |
| | Standalone | Hepatitis B Surface Antigen Prenatal, Serum | ✓ | HBAGP | 87340 | http://www.mayomedicallaboratories.com/test-catalog/Overview/86185 |
| Quest Diagnostics | Panel | Obstetric Panel | ✓ | 20210 | 80055 | http://www.questdiagnostics.com/testcenter/BUOrderInfo.action?tc=20210&labCode=MIA |
| | Standalone | N/A | N/A | N/A | N/A | |

*When an HBsAg test result is reactive, laboratories may automatically perform a confirmatory test without additional provider order.

**This CPT code corresponds only to the HBsAg screening component of this laboratory panel; additional CPT codes might be associated with other component tests in this laboratory panel.

Notes: CDC recommends healthcare providers use prenatal HBsAg tests (vs. non-specific tests) for pregnant women, which allows for reporting of positive results along with pregnancy status to public health jurisdictions. Refer all HBsAg positive pregnant women to Perinatal Hepatitis B Prevention Program coordinators for case management of mother and infant:
<http://www.cdc.gov/vaccines/vpd-vac/hepb/perinatal-contacts.htm>.

Laboratories reserve the right to add, modify, or stop performing tests at any time – providers should review any test notifications from laboratories for changes.




U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



The American College of Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

www.cdc.gov/hepatitis

Screening and Referral Algorithm for Hepatitis B Virus (HBV) Infection among Pregnant Women



*High risk for HBV infection includes: household or sexual contacts of HBsAg-positive persons; injection drug use; more than one sex partner during the past six months; evaluation or treatment for a sexually transmitted disease; HIV infection, chronic liver disease, or end-stage renal disease; and international travel to regions with HBsAg prevalence of $\geq 2\%$.

Adapted with permission from the Hepatitis B Foundation. Original publication: Apuzzio J, Block J, Cullison S, et al. Chronic Hepatitis B in pregnancy: A workshop consensus statement on screening, evaluation, and management, part 2. *The Female Patient*. 2012; 37(5):30-34



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



The American College of Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

www.cdc.gov/hepatitis

March 2015